

Patient **Able, Denny**
UNIT #: **1234567**

DOB: **09/02/1941**

AUDIOLOGIST: **John Doe, Au.D.**

DATE: **01/06/2005**

AUDIOMETER: **GSI 61**

TEST METHOD: **Conventional**

RELIABILITY: **Good**

Air Conduction	Right	Left
Unmasked	O	X
Masked	Δ	□
Bone Conduction		
Unmasked	<	>
Masked	◁	▷
Sound Field		
Unaided - Aided	S	A
Comfort Level		
Maximum	MC	MC
Uncomfortable	UC	UC
Reflexes		
Contra	↖	↗
Ipsi	↘	↙
Commented	C / P / V	

C = see comments, P = Pressure
V = Vibrotactile, N = Narrow Band Noise
FM = Frequency Modulation

Tympanometry

	Right	Left
Probe Tone (Hz)	226	226
Ear Canal Volume	1.30	1.60
Peak Admittance (ml)	0.70	1.40
Peak pressure (daPa)	15	-45
Curve Type	A	As

Acoustic Reflex Results

	Right		Left		Right	Left
	Ipsi	Contra	Ipsi	Contra	Reflex Decay	
500	80	Abs	Abs			Abnrml
1000	CNT	95	UdB		Nrml	
2000						
4000	70	105	55	50		

Abs- Absent CNT- Could Not Test UdB- Undefined decibel level

* = masked, NR = No Response

- ☐ Supraaural Earphones ☐ Inserts ☐ Soundfield
☐ Pure Tone ☒ Warble ☐ Narrow Band Noise

SPEECH AUDIOMETRY (Results are in dB)

	REC	MLV	UNAIDED			AIDED		
			RIGHT	LEFT	SOUNDFIELD	RIGHT	LEFT	[Phonic Bivale] BINAURAL
SAT [ba/sh/s]		X	25	35	45	55	65	75
SRT			20	30		50	60	70
Maryland CNC / LV		X	* 20 % at 30					
Maryland CNC / REC	X		50 % at 50					
NU-6 / LV		X	90 % at 70					
NU-6 / REC	X		40 % at 90					
W-22 / LV		X	25 % at 10					
W-22 / REC	X		50 % at 25					
WD 25 / LV		X	1 % at 2					
WD 25 / REC	X		3 % at 4					

COMMENTS

This is a test of the text measurments,
Wound care instructions: 1) Keep steri-strips dry for first 24 hours, then you can let water and soap run over them. Just gently pat them dry afterwards. 2) The steri-strips should stay on for a week. After that you can remove them whenever you want. They will gradually peel off and typically fall off by themselves the second week after surgery. You can just trim the edges as they come up during the first week.

EXAMINER SIGNATURE